

# Multiple Regression Analysis - Inference

Caio Vigo

**The University of Kansas**  
Department of Economics

Spring 2020

These slides were based on *Introductory Econometrics* by Jeffrey M. Wooldridge (2015)

## Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

### 1 Motivation

### 2 Sampling Distributions of the OLS Estimators

### 3 Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

### 4 Confidence Intervals

### 5 Testing Multiple Exclusion Restrictions

$R$ -Squared Form of the  $F$  Statistic

The  $F$  Statistic for Overall Significance of a Regression

## Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

**Goal:** We want to test hypothesis about the parameters  $\beta_j$  in the population regression model.

We want to know whether the true parameter  $\beta_j = \text{some value (your hypothesis)}$ .

- In order to do that, we will need to add a final assumption **MLR.6**. We will obtain the **Classical Linear Model (CLM)**

## Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

**MLR.1:**  $y = \beta_0 + \beta_1x_1 + \beta_2x_2 + \dots + \beta_kx_k + u$

**MLR.2:** random sampling from the population

**MLR.3:** no perfect collinearity in the sample

**MLR.4:**  $E(u|x_1, \dots, x_k) = E(u) = 0$  (exogenous explanatory variables)

**MLR.5:**  $Var(u|x_1, \dots, x_k) = Var(u) = \sigma^2$  (homoskedasticity)

**MLR.1 - MLR.4:** Needed for unbiasedness of OLS:

$$E(\hat{\beta}_j) = \beta_j$$

**MLR.1 - MLR.5:** Needed to compute  $Var(\hat{\beta}_j)$ :

$$Var(\hat{\beta}_j) = \frac{\sigma^2}{SST_j(1 - R_j^2)}$$

$$\hat{\sigma}^2 = \frac{SSR}{(n - k - 1)}$$

and for efficiency of OLS  $\Rightarrow$  **BLUE**.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- ① Motivation
- ② Sampling Distributions of the OLS Estimators
- ③ Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- ④ Confidence Intervals
- ⑤ Testing Multiple Exclusion Restrictions
  - $R$ -Squared Form of the  $F$  Statistic
  - The  $F$  Statistic for Overall Significance of a Regression

- Now we need to know the full sampling distribution of the  $\hat{\beta}_j$ .
- The **Gauss-Markov assumptions** don't tell us anything about these distributions.
- Based on our models, (conditional on  $\{(x_{i1}, \dots, x_{ik}) : i = 1, \dots, n\}$ ) we need to have  $dist(\hat{\beta}_j) = f(dist(u))$ , i.e.,

$$\hat{\beta}_j \sim pdf(u)$$

- That's why we need one more assumption.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## MRL.6 (Normality)

The population error  $u$  is independent of the explanatory variables  $(x_1, \dots, x_k)$  and is normally distributed with mean zero and variance  $\sigma^2$ :

$$u \sim \text{Normal}(0, \sigma^2)$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

**MLR.1 - MLR.4**  $\rightarrow$  unbiasedness of OLS

**Gauss-Markov assumptions:** **MLR.1 - MLR.4** + **MLR.5** (homoskedastic errors)

**Classical Linear Model (CLM):** **Gauss-Markov** + **MLR.6** (Normally distributed errors)



- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

$$u \sim \text{Normal}(0, \sigma^2)$$

- Strongest assumption.
- **MLR.6** implies  $\Rightarrow$  zero conditional mean (**MLR.4**) and homoskedasticity (**MLR.5**)
- Now we have full independence between  $u$  and  $(x_1, x_2, \dots, x_k)$  (*not just mean and variance independence*)
- Reason to call  $x_j$  **independent variables**.
- Recall the Normal distribution properties (see slides for **Appendix B**).

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$


Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals


Testing Multiple Exclusion

Figure: Distribution of  $u$ :  $u \sim N(0, \sigma^2)$



- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

Figure:  $f(y|x)$  with homoskedastic normal errors, i.e.,  $u \sim N(0, \sigma^2)$



- Property of a **Normal distribution**: if  $W \sim Normal$  then  $a + bW \sim Normal$  for constants  $a$  and  $b$ .

- What we are saying is that for normal r.v.s, any linear combination of them is also normally distributed.
- Because the  $u_i$  are *independent and identically distributed (iid)* as  $Normal(0, \sigma^2)$

$$\hat{\beta}_j = \beta_j + \sum_{i=1}^n w_{ij}u_i \sim Normal(\beta_j, Var(\hat{\beta}_j))$$

- Then we can apply the Central Limit Theorem.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

## Theorem: Normal Sampling Distributions

Under the **CLM** assumptions, conditional on the sample outcomes of the explanatory variables,

$$\hat{\beta}_j \sim \text{Normal}(\beta_j, \text{Var}(\hat{\beta}_j))$$

and so

$$\frac{\hat{\beta}_j - \beta_j}{\text{sd}(\hat{\beta}_j)} \sim \text{Normal}(0, 1)$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- ① Motivation
- ② Sampling Distributions of the OLS Estimators
- ③ Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- ④ Confidence Intervals
- ⑤ Testing Multiple Exclusion Restrictions
  - $R$ -Squared Form of the  $F$  Statistic
  - The  $F$  Statistic for Overall Significance of a Regression

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## Theorem: $t$ Distribution for Standardized Estimators

Under the **CLM** assumptions,

$$\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \sim t_{n-k-1} = t_{df}$$

where  $k + 1$  is the number of unknown parameter in the population model, and  $n - k - 1$  is the degrees of freedom (df).

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- Compare the ratios of the **previous 2** theorems. What is the difference?
- What is the difference between  $sd(\hat{\beta}_j)$  and  $se(\hat{\beta}_j)$ ?
- Recall the  $t$  distribution properties (see slides for **Appendix B**).



Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- The  $t$  distribution also has a bell shape, but is more spread out than the  $Normal(0, 1)$ .


- As  $df \rightarrow \infty$ ,

$$t_{df} \rightarrow Normal(0, 1)$$

- The difference is practically small for  $df > 120$ .
- See a  $t$  table.
- The next graph plots a standard normal pdf against a  $t_6$  pdf.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter**
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

Figure: The pdfs of a standard normal and a  $t_6$



- We use the result on the  $t$  distribution to test the null hypothesis that  $x_j$  has no partial effect on  $y$ :

$$H_0 : \beta_j = 0$$

$$lwage = \beta_0 + \beta_1 educ + \beta_2 exper + \beta_3 tenure + u$$

$$H_0 : \beta_2 = 0$$

- **Interpretation of what we are doing:** Once we control for education and time on the current job (*tenure*), total workforce experience has no affect on  $lwage = \log(wage)$ .

- To test

$$H_0 : \beta_j = 0$$

we use the **t statistic** (or **t ratio**),

$$t_{\hat{\beta}_j} = \frac{\hat{\beta}_j}{se(\hat{\beta}_j)}$$

- In virtually all cases  $\hat{\beta}_j$  is not exactly equal to zero.
- When we use  $t_{\hat{\beta}_j}$ , we are measuring how far  $\hat{\beta}_j$  is from zero *relative to its standard error*.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- First consider the alternative

$$H_1 : \beta_j > 0$$

which means the null is effectively

$$H_0 : \beta_j \leq 0$$

- Using a positive one-sided alternative, if we reject  $\beta_j = 0$ , then we reject any  $\beta_j < 0$ , too.
- We often just state  $H_0 : \beta_j = 0$  and act like we do not care about negative values.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- Because  $se(\hat{\beta}_j) > 0$ ,  $t_{\hat{\beta}_j}$  always has the same sign as  $\hat{\beta}_j$ .
- If the estimated coefficient  $\hat{\beta}_j$  is negative, it provides no evidence against  $H_0$  in favor of  $H_1 : \beta_j > 0$ .
- If  $\hat{\beta}_j$  is positive, the question is: How big does  $t_{\hat{\beta}_j} = \hat{\beta}_j / se(\hat{\beta}_j)$  have to be before we conclude  $H_0$  is “unlikely”?
- Let's review the Error Types in Statistics.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- Consider the following example:

$H_0$  : Not pregnant

$H_1$  : Pregnant

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$


Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

**Type I error**  
(false positive)



**Type II error**  
(false negative)





- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

|               |           | Reality<br>H0 is actually:             |                                       |
|---------------|-----------|--|---------------------------------------|
|               |           | False                                  | True                                  |
| Study Finding | Reject H0 | <b>True Positive</b><br>(Power)        | False Positive<br><b>Type I Error</b> |
|               | Accept H0 | False Negative<br><b>Type II Error</b> | <b>True Negative</b>                  |

# Testing Against One-Sided Alternatives

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

1. Choose a null hypothesis:  $H_0 : \beta_j = 0$  (or  $H_0 : \beta_j \leq 0$ )

2. Choose an alternative hypothesis:  $H_1 : \beta_j > 0$

3. Choose a **significance level**  $\alpha$  (or simply **level**, or **size**) for the test.

That is, the probability of rejecting the null hypothesis when it is in fact true. (Type I Error).

Suppose we use 5%, so the probability of committing a Type I error is .05.

4. Obtain the critical value,  $c > 0$ , so that the **rejection rule**

$$t_{\hat{\beta}_j} > c$$

leads to a 5% level test.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- The key is that, *under the null hypothesis*,


$$t_{\hat{\beta}_j} \sim t_{n-k-1} = t_{df}$$

and this is what we use to obtain the critical value,  $c$ .

- Suppose  $df = 28$  and we use a 5% test.
- Find the **critical value** in a t-table. (table).

# Testing Against One-Sided Alternatives

Numbers in each row of the table are values on a  $t$ -distribution with ( $df$ ) degrees of freedom for selected right-tail (greater-than) probabilities ( $p$ ).



| $df/p$    | 0.40     | 0.25     | 0.10     | 0.05     | 0.025    | 0.01     | 0.005    | 0.0005   |
|-----------|----------|----------|----------|----------|----------|----------|----------|----------|
| <b>1</b>  | 0.324920 | 1.000000 | 3.077684 | 6.313752 | 12.70620 | 31.82052 | 63.65674 | 636.6192 |
| <b>2</b>  | 0.288675 | 0.816497 | 1.885618 | 2.919986 | 4.30265  | 6.96456  | 9.92484  | 31.5991  |
| <b>3</b>  | 0.276671 | 0.764892 | 1.637744 | 2.353363 | 3.18245  | 4.54070  | 5.84091  | 12.9240  |
|           |          |          |          |          |          |          |          |          |
| <b>25</b> | 0.256060 | 0.684430 | 1.316345 | 1.708141 | 2.05954  | 2.48511  | 2.78744  | 3.7251   |
| <b>26</b> | 0.255955 | 0.684043 | 1.314972 | 1.705618 | 2.05553  | 2.47863  | 2.77871  | 3.7066   |
| <b>27</b> | 0.255858 | 0.683685 | 1.313703 | 1.703288 | 2.05183  | 2.47266  | 2.77068  | 3.6896   |
| <b>28</b> | 0.255768 | 0.683353 | 1.312527 | 1.701131 | 2.04841  | 2.46714  | 2.76326  | 3.6739   |
| <b>29</b> | 0.255684 | 0.683044 | 1.311434 | 1.699127 | 2.04523  | 2.46202  | 2.75639  | 3.6594   |
| <b>30</b> | 0.255605 | 0.682756 | 1.310415 | 1.697261 | 2.04227  | 2.45726  | 2.75000  | 3.6460   |
| <b>z</b>  | 0.253347 | 0.674490 | 1.281552 | 1.644854 | 1.95996  | 2.32635  | 2.57583  | 3.2905   |
| <b>CI</b> | ————     | ————     | 80%      | 90%      | 95%      | 98%      | 99%      | 99.9%    |

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests


Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- The critical value is  $c = 1.701$  for 5% significance level (one-sided test).
- The following picture shows that we are conducting a **one-tailed test** (and it is these entries that should be used in the table).

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion



- So, with  $df = 28$ , the rejection rule for  $H_0 : \beta_j = 0$  against  $H_1 : \beta_j > 0$ , at the 5% level, is

$$t_{\hat{\beta}_j} > 1.701$$


We need a  $t$  statistic greater than 1.701 to conclude there is enough evidence against  $H_0$ .

- If  $t_{\hat{\beta}_j} \leq 1.701$ , we fail to reject  $H_0$  against  $H_1$  at the 5% significance level.

# Testing Against One-Sided Alternatives

- Suppose  $df = 28$ , but we want to carry out the test at a different significance level (often 10% level or the 1% level).

Numbers in each row of the table are values on a  $t$ -distribution with ( $df$ ) degrees of freedom for selected right-tail (greater-than) probabilities ( $p$ ).



| $df/p$ | 0.40     | 0.25     | 0.10     | 0.05     | 0.025    | 0.01     | 0.005    | 0.0005   |
|--------|----------|----------|----------|----------|----------|----------|----------|----------|
| 1      | 0.324920 | 1.000000 | 3.077684 | 6.313752 | 12.70620 | 31.82052 | 63.65674 | 636.6192 |
| 2      | 0.288675 | 0.816497 | 1.885618 | 2.919986 | 4.30265  | 6.96456  | 9.92484  | 31.5991  |
| 3      | 0.276671 | 0.764892 | 1.637744 | 2.353363 | 3.18245  | 4.54070  | 5.84091  | 12.9240  |
|        |          |          |          |          |          |          |          |          |
| 25     | 0.256060 | 0.684430 | 1.316345 | 1.708141 | 2.05954  | 2.48511  | 2.78744  | 3.7251   |
| 26     | 0.255955 | 0.684043 | 1.314972 | 1.705618 | 2.05553  | 2.47863  | 2.77871  | 3.7066   |
| 27     | 0.255858 | 0.683685 | 1.313703 | 1.703288 | 2.05183  | 2.47266  | 2.77068  | 3.6896   |
| 28     | 0.255768 | 0.683353 | 1.312527 | 1.701131 | 2.04841  | 2.46714  | 2.76326  | 3.6739   |
| 29     | 0.255684 | 0.683044 | 1.311434 | 1.699127 | 2.04523  | 2.46202  | 2.75639  | 3.6594   |
| 30     | 0.255605 | 0.682756 | 1.310415 | 1.697261 | 2.04227  | 2.45726  | 2.75000  | 3.6460   |
| $z$    | 0.253347 | 0.674490 | 1.281552 | 1.644854 | 1.95996  | 2.32635  | 2.57583  | 3.2905   |
| CI     | ————     | ————     | 80%      | 90%      | 95%      | 98%      | 99%      | 99.9%    |

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion



- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- Thus, if  $df = 28$ , below are the critical values for the following significance levels: 10% level, 5% and 1% level.

$$c_{.10} = 1.313$$

$$c_{.05} = 1.701$$

$$c_{.01} = 2.467$$

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

If we want to reduce the probability of Type I error, we must increase the critical value (so we reject the null less often).

- If we reject at, say, the 1% level, then we must also reject at any larger level.
- If we fail to reject at, say, the 10% level – so that  $t_{\hat{\beta}_j} \leq 1.313$  – then we will fail to reject at any smaller level.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- With large sample sizes – certain when  $df > 120$  – we can use critical values from the standard normal distribution.

$$c_{.10} = 1.282$$

$$c_{.05} = 1.645$$

$$c_{.01} = 2.326$$

which we can round to 1.28, 1.65, and 2.36, respectively. The value 1.65 is especially common for a one-tailed test.

- Recall our **wage** model example:

$$\log(\text{wage}) = \beta_0 + \beta_1 \text{educ} + \beta_2 \text{exper} + \beta_3 \text{tenure} + u$$

- First, let's label the parameters with the variable names:  $\beta_{\text{educ}}$ ,  $\beta_{\text{exper}}$ , and  $\beta_{\text{tenure}}$
- We would like to test:

$$H_0 : \beta_{\text{exper}} = 0$$

**Interpretation:** We are testing if workforce experience has no effect on a wage once education and tenure have been accounted for.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

```

=====
                                Dependent variable:
                                -----
                                lwage
-----
educ                            0.092***
                                (0.007)
exper                            0.004**
                                (0.002)
tenure                           0.022***
                                (0.003)
Constant                         0.284***
                                (0.104)
-----
Observations                     526
R2                               0.316
Adjusted R2                      0.312
Residual Std. Error              0.441 (df = 522)
F Statistic                      80.391*** (df = 3; 522)
=====
Note:                            *p<0.1; **p<0.05; ***p<0.01
  
```

- What is the  $t_{exper}$ ?

$$t_{exper} = \frac{0.004}{0.002} = 2.00$$

- Now what do you do with this number?
- How many  $df$  do we have?
- Which table could I use?
- Using a standard normal table: the one-sided critical value at the 5% level, 1.645.

## Statistical Significance X Economic Importance/Interpretation

- So “ $\hat{\beta}_{exper}$  is **statistically significant**” at 5% level significance level (one-sided test).
- The estimated effect of *exper*, which is its **economic importance** should be interpreted as: another year of experience, holding *educ* and *tenure* fixed, is estimated to be worth about 0.4%.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- For the negative one-sided alternative,

$$H_0 : \beta_j \geq 0$$

$$H_1 : \beta_j < 0$$

we use a symmetric rule. But the rejection rule is

$$t_{\hat{\beta}_j} < -c$$


where  $c$  is chosen in the same way as in the positive case.



# Testing Against One-Sided Alternatives

- With  $df = 28$  and we want to test at a 5% significance level, what is the critical value?

Numbers in each row of the table are values on a  $t$ -distribution with ( $df$ ) degrees of freedom for selected right-tail (greater-than) probabilities ( $p$ ).



| df/p | 0.40     | 0.25     | 0.10     | 0.05     | 0.025    | 0.01     | 0.005    | 0.0005   |
|------|----------|----------|----------|----------|----------|----------|----------|----------|
| 1    | 0.324920 | 1.000000 | 3.077684 | 6.313752 | 12.70620 | 31.82052 | 63.65674 | 636.6192 |
| 2    | 0.288675 | 0.816497 | 1.885618 | 2.919986 | 4.30265  | 6.96456  | 9.92484  | 31.5991  |
| 3    | 0.276671 | 0.764892 | 1.637744 | 2.353363 | 3.18245  | 4.54070  | 5.84091  | 12.9240  |
|      |          |          |          |          |          |          |          |          |
| 25   | 0.256060 | 0.684430 | 1.316345 | 1.708141 | 2.05954  | 2.48511  | 2.78744  | 3.7251   |
| 26   | 0.255955 | 0.684043 | 1.314972 | 1.705618 | 2.05553  | 2.47863  | 2.77871  | 3.7066   |
| 27   | 0.255858 | 0.683685 | 1.313703 | 1.703288 | 2.05183  | 2.47266  | 2.77068  | 3.6896   |
| 28   | 0.255768 | 0.683353 | 1.312527 | 1.701131 | 2.04841  | 2.46714  | 2.76326  | 3.6739   |
| 29   | 0.255684 | 0.683044 | 1.311434 | 1.699127 | 2.04523  | 2.46202  | 2.75639  | 3.6594   |
| 30   | 0.255605 | 0.682756 | 1.310415 | 1.697261 | 2.04227  | 2.45726  | 2.75000  | 3.6460   |
| z    | 0.253347 | 0.674490 | 1.281552 | 1.644854 | 1.95996  | 2.32635  | 2.57583  | 3.2905   |
| CI   | ————     | ————     | 80%      | 90%      | 95%      | 98%      | 99%      | 99.9%    |

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals


Testing Multiple Exclusion

**Intuition:** We must see a significantly negative value for the  $t$  statistic to reject the null hypothesis in favor of the alternative hypothesis.

- With  $df = 28$  and a 5% test, the critical value is  $c = -1.701$ , so the rejection rule is

$$t_{\hat{\beta}_j} < -1.701$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives**
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion



## Reminder about Testing

- Our hypotheses involve the unknown population values,  $\beta_j$ .
- If in a our set of data we obtain, say,  $\hat{\beta}_j = 2.75$ , we do not write the null hypothesis as

$$H_0 : 2.75 = 0$$

(which is obviously false).

- Nor do we write

$$H_0 : \hat{\beta}_j = 0$$

(which is also false except in the very rare case that our estimate is exactly zero).

- We do not test hypotheses about the estimate! We know what it is once we collect the sample. We hypothesize about the unknown population value,  $\beta_j$ .

## Testing Against Two-Sided Alternatives

- Sometimes we do not know ahead of time whether a variable definitely has a positive effect or a negative effect.
- So, in this case the hypothesis should be written as:

$$H_0 : \beta_j = 0$$

$$H_1 : \beta_j \neq 0$$

- Testing against the **two-sided alternative** is usually the default. It prevents us from looking at the regression results and then deciding on the alternative.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

**Testing Against Two-Sided Alternatives**

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion


- Now we reject if  $\hat{\beta}_j$  is sufficiently large in magnitude, either positive or negative. We again use the  $t$  statistic  $t_{\hat{\beta}_j} = \hat{\beta}_j / se(\hat{\beta}_j)$ , but now the rejection rule is

## Two-tailed test

$$|t_{\hat{\beta}_j}| > c$$

- For example, if we use a 5% level test and  $df = 25$ , the two-tailed cv is 2.06. The two-tailed cv is, in this case, the 97.5 percentile in the  $t_{25}$  distribution. (Compare the one-tailed cv, about 1.71, the 95<sup>th</sup> percentile in the  $t_{25}$  distribution).


- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives**
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion





# Testing Against Two-Sided Alternatives

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives**
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion



|     | 0.00   | 0.01   | 0.02   | 0.03   | 0.04   | 0.05   | 0.06   | 0.07   | 0.08   | 0.09   |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0.0 | 0.5000 | 0.5040 | 0.5080 | 0.5120 | 0.5160 | 0.5199 | 0.5239 | 0.5279 | 0.5319 | 0.5359 |
| 0.1 | 0.5398 | 0.5438 | 0.5478 | 0.5517 | 0.5557 | 0.5596 | 0.5636 | 0.5675 | 0.5714 | 0.5753 |
| 0.2 | 0.5793 | 0.5832 | 0.5871 | 0.5910 | 0.5948 | 0.5987 | 0.6026 | 0.6064 | 0.6103 | 0.6141 |
| ⋮   | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      | ⋮      |
| 1.5 | 0.9332 | 0.9345 | 0.9357 | 0.9370 | 0.9382 | 0.9394 | 0.9406 | 0.9418 | 0.9429 | 0.9441 |
| 1.6 | 0.9452 | 0.9463 | 0.9474 | 0.9484 | 0.9495 | 0.9505 | 0.9515 | 0.9525 | 0.9535 | 0.9545 |
| 1.7 | 0.9554 | 0.9564 | 0.9573 | 0.9582 | 0.9591 | 0.9599 | 0.9608 | 0.9616 | 0.9625 | 0.9633 |
| 1.8 | 0.9641 | 0.9649 | 0.9656 | 0.9664 | 0.9671 | 0.9678 | 0.9686 | 0.9693 | 0.9699 | 0.9706 |
| 1.9 | 0.9713 | 0.9719 | 0.9726 | 0.9732 | 0.9738 | 0.9744 | 0.9750 | 0.9756 | 0.9761 | 0.9767 |
| 2.0 | 0.9772 | 0.9778 | 0.9783 | 0.9788 | 0.9793 | 0.9798 | 0.9803 | 0.9808 | 0.9812 | 0.9817 |
| 2.1 | 0.9821 | 0.9826 | 0.9830 | 0.9834 | 0.9838 | 0.9842 | 0.9846 | 0.9850 | 0.9854 | 0.9857 |
| 2.2 | 0.9861 | 0.9864 | 0.9868 | 0.9871 | 0.9875 | 0.9878 | 0.9881 | 0.9884 | 0.9887 | 0.9890 |
| 2.3 | 0.9893 | 0.9896 | 0.9898 | 0.9901 | 0.9904 | 0.9906 | 0.9909 | 0.9911 | 0.9913 | 0.9916 |
| 2.4 | 0.9918 | 0.9920 | 0.9922 | 0.9925 | 0.9927 | 0.9929 | 0.9931 | 0.9932 | 0.9934 | 0.9936 |
| 2.5 | 0.9938 | 0.9940 | 0.9941 | 0.9943 | 0.9945 | 0.9946 | 0.9948 | 0.9949 | 0.9951 | 0.9952 |
| 2.6 | 0.9953 | 0.9955 | 0.9956 | 0.9957 | 0.9959 | 0.9960 | 0.9961 | 0.9962 | 0.9963 | 0.9964 |
| 2.7 | 0.9965 | 0.9966 | 0.9967 | 0.9968 | 0.9969 | 0.9970 | 0.9971 | 0.9972 | 0.9973 | 0.9974 |
| 2.8 | 0.9974 | 0.9975 | 0.9976 | 0.9977 | 0.9977 | 0.9978 | 0.9979 | 0.9979 | 0.9980 | 0.9981 |
| 2.9 | 0.9981 | 0.9982 | 0.9982 | 0.9983 | 0.9984 | 0.9984 | 0.9985 | 0.9985 | 0.9986 | 0.9986 |
| 3.0 | 0.9987 | 0.9987 | 0.9987 | 0.9988 | 0.9988 | 0.9989 | 0.9989 | 0.9989 | 0.9990 | 0.9990 |

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

```

=====
                                Dependent variable:
                                -----
                                lwage
-----
educ                            0.092***
                                (0.007)

exper                            0.004**
                                (0.002)

tenure                           0.022***
                                (0.003)

Constant                         0.284***
                                (0.104)

-----
Observations                     526
R2                               0.316
Adjusted R2                      0.312
Residual Std. Error             0.441 (df = 522)
F Statistic                     80.391*** (df = 3; 522)
=====
Note:                            *p<0.1; **p<0.05; ***p<0.01
  
```

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- When we reject  $H_0 : \beta_j = 0$  against  $H_1 : \beta_j \neq 0$ , we often say that  $\hat{\beta}_j$  is **statistically different from zero** and usually mention a significance level.

As in the one-sided case, we also say  $\hat{\beta}_j$  is **statistically significant** when we can reject  $H_0 : \beta_j = 0$ .

Motivation

Sampling  
Distributions  
of the OLS  
Estimators

Testing  
Hypotheses  
About a Single  
Population  
Parameter

Testing Against  
One-Sided  
Alternatives

Testing Against  
Two-Sided  
Alternatives

Testing Other  
Hypotheses about  
the  $\beta_j$

Computing  $p$ -Values  
for  $t$  Tests

Practical (Economic)  
versus Statistical  
Significance

Confidence  
Intervals

Testing  
Multiple  
Exclusion

- Testing the null  $H_0 : \beta_j = 0$  is the standard practice.
- **R**, Stata, EViews and all the other regression packages automatically report the  $t$  statistic for **this hypothesis** (i.e., two-sided test).

- What if we want to test a different null value? For example, in a constant-elasticity consumption function,

$$\log(\text{cons}) = \beta_0 + \beta_1 \log(\text{inc}) + \beta_2 \text{famsize} + \beta_3 \text{pareduc} + u$$

we might want to test

$$H_0 : \beta_1 = 1$$

which means an income elasticity equal to one. (We can be pretty sure that  $\beta_1 > 0$ .)

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## Important observation

$$t_{\hat{\beta}_j} = \frac{\hat{\beta}_j}{se(\hat{\beta}_j)}$$

is *only* for  $H_0 : \beta_j = 0$ .

# Testing Other Hypotheses about the $\beta_j$

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- More generally, suppose the null is

$$H_0 : \beta_j = a_j$$

where we specify the value  $a_j$

- It is easy to extend the  $t$  statistic:

$$t = \frac{(\hat{\beta}_j - a_j)}{se(\hat{\beta}_j)}$$

The  $t$  statistic just measures how far our estimate,  $\hat{\beta}_j$ , is from the hypothesized value,  $a_j$ , relative to  $se(\hat{\beta}_j)$ .

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## General expression for general $t$ testing

$$t = \frac{(\textit{estimate} - \textit{hypothesized value})}{\textit{standard error}}$$

- The alternative can be one-sided or two-sided.
- We choose critical values in exactly the same way as before.



- The language needs to be suitably modified. If, for example,

$$H_0 : \beta_j = 1$$

$$H_1 : \beta_j \neq 1$$

is rejected at the 5% level, we say “ $\hat{\beta}_j$  is statistically different from one at the 5% level.” Otherwise,  $\hat{\beta}_j$  is “not statistically different from one.” If the alternative is  $H_1 : \beta_j > 1$ , then “ $\hat{\beta}_j$  is statistically greater than one at the 5% level.”

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

**Example:** Crime, police officers and enrollment on college campuses  
 Let's do the following hypothesis test:

$$\log(\textit{crime}) = \beta_0 + \beta_1 \textit{police} + \beta_2 \log(\textit{enroll}) + u$$

$$H_0 : \beta_1 = 1$$

$$H_1 : \beta_1 > 1$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

```

=====
                                Dependent variable:
                                -----
                                log(crime)
-----
police                            0.0240***
                                   (0.0073)

log(enroll)                        0.9767***
                                   (0.1373)

Constant                          -4.3758***
                                   (1.1990)

-----
Observations                        97
R2                                  0.6277
Adjusted R2                         0.6198
Residual Std. Error                0.8516 (df = 94)
F Statistic                        79.2389*** (df = 2; 94)
=====
Note:                               *p<0.1; **p<0.05; ***p<0.01
  
```

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- The traditional approach to testing, where we choose a significance level ahead of time, has a component of **arbitrariness**.
- Different researchers prefer different significance levels (10%, 5%, 1%).
- Committing to a significance level ahead of time can hide useful information about the outcome of hypothesis test.
- **Example:** (On white board)

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- Rather than have to specify a level ahead of time, or discuss different traditional significance levels (10%, 5%, 1%), it is better to answer the following question:

**Intuition:** Given the observed value of the  $t$  statistic, what is the *smallest* significance level at which I can reject  $H_0$ ?

- The smallest level at which the null can be rejected is known as the  **$p$ -value** of a test.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## $p$ -value

For  $t$  testing against a two-sided alternative,

$$p\text{-value} = P(|T| > |t|)$$

where  $t$  is the value of the  $t$  statistic and  $T$  is a random variable with the  $t_{df}$  distribution.

- The  $p$ -value is a probability, so it is between zero and one.

Motivation

Sampling  
Distributions  
of the OLS  
Estimators

Testing  
Hypotheses  
About a Single  
Population  
Parameter

Testing Against  
One-Sided  
Alternatives

Testing Against  
Two-Sided  
Alternatives

Testing Other  
Hypotheses about  
the  $\beta_j$

**Computing  $p$ -Values  
for  $t$  Tests**

Practical (Economic)  
versus Statistical  
Significance

Confidence  
Intervals

Testing  
Multiple  
Exclusion

One way to think about the  $p$ -values is that it uses the observed statistic as the critical value, and then finds the significance level of the test using that critical value.

- Usually we just report  $p$ -values for two-sided alternatives.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## Mnemonic Device

*Small*  $p$ -values are evidence *against* the null hypothesis.

*Large*  $p$ -values provide little evidence *against* the null hypothesis.

**Intuition:**  $p$ -value is the probability of observing a statistic as extreme as we did if the null hypothesis is true.



Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- If  $p$ -value = .50, then there is a 50% chance of observing a  $t$  as large as we did (in absolute value). This is not enough evidence against  $H_0$ .
- If  $p$ -value = .001, then the chance of seeing a  $t$  statistic as extreme as we did is .1%.
- We can conclude that we got a very rare sample (*unlikely!*) or that the null hypothesis is very likely false.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests**
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- From

$$p\text{-value} = P(|T| > |t|)$$

we see that as  $|t|$  increases the  $p$ -value decreases.

Large absolute  $t$  statistics are **associated** with small  $p$ -values.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

## Example:


- Suppose  $df = 40$  and, from our data, we obtain  $t = 1.85$  or  $t = -1.85$ . Then

$$p\text{-value} = P(|T| > 1.85) = 2P(T > 1.85) = 2(.0359) = .0718$$

where  $T \sim t_{40}$ .

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

Figure:  $t$  distribution with 40 degrees of freedom



- Given  $p$ -value, we can carry out a test at any significance level. If  $\alpha$  is the chosen level, then

Reject  $H_0$  if  $p\text{-value} < \alpha$

## Example

Suppose we obtained  $p\text{-value} = .0718$ . This means that we reject  $H_0$  at the 10% level but not the 5% level. We reject at 8% but not at 7%.

Motivation

Sampling  
Distributions  
of the OLS  
Estimators

Testing  
Hypotheses  
About a Single  
Population  
Parameter

Testing Against  
One-Sided  
Alternatives

Testing Against  
Two-Sided  
Alternatives

Testing Other  
Hypotheses about  
the  $\beta_j$

Computing  $p$ -Values  
for  $t$  Tests

Practical (Economic)  
versus Statistical  
Significance

Confidence  
Intervals

Testing  
Multiple  
Exclusion

- $t$  testing is purely about *statistical significance*.
- It does not directly speak to the issue of whether a variable has a practically, or economically large effect.

**Practical (Economic) Significance** depends on the size (and sign) of  $\hat{\beta}_j$ .

**Statistical Significance** depends on  $t_{\hat{\beta}_j}$ .

# Practical (Economic) versus Statistical Significance

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

It is possible estimate **practically large effects** but have the estimates so imprecise that they are **statistically insignificant**.

Common with small data sets (but not only small data sets).

**X**

It is possible to get estimates that are **statistically significant** (often with very small  $p$ -values) but are **not practically large**.

Common with very large data sets.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- ① Motivation
- ② Sampling Distributions of the OLS Estimators
- ③ Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- ④ Confidence Intervals
- ⑤ Testing Multiple Exclusion Restrictions
  - $R$ -Squared Form of the  $F$  Statistic
  - The  $F$  Statistic for Overall Significance of a Regression



Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

**Confidence Intervals**

Testing Multiple Exclusion

- **Under the CLM assumptions**, rather than just testing hypotheses about parameters it is also useful to construct **confidence intervals** (also know as **interval estimates**).

**Intuition:** If you could obtain several random samples data, the **confidence interval** tells you that, for a 95% CI, your true  $\beta_j$  will lie in this interval  $[\beta_j^{lower}, \beta_j^{upper}]$  for 95% of the samples.

- We will construct CIs of the form

$$\hat{\beta}_j \pm c \cdot se(\hat{\beta}_j)$$

where  $c > 0$  is chosen based on the **confidence level**.

- We will use a 95% confidence level, in which case  $c$  comes from the 97.5 percentile in the  $t_{df}$  distribution.
- Therefore,  $c$  is the 5% critical value against a two-sided alternative.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

**Confidence Intervals**

Testing Multiple Exclusion

## Example

- For,  $df \geq 120$ , the 95% CI is:

$$\hat{\beta}_j \pm 1.96 \cdot se(\hat{\beta}_j) \text{ or } \left[ \hat{\beta}_j - 1.96 \cdot se(\hat{\beta}_j), \hat{\beta}_j + 1.96 \cdot se(\hat{\beta}_j) \right]$$

- For small  $df$ , the exact percentiles should be obtained from a  $t$  table.

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals**
- Testing Multiple Exclusion

Find the 95% CI for the parameters from the following regression:

```

=====
                                Dependent variable:
-----
                                lwage
-----
educ                            0.092***
                                (0.007)

exper                            0.004**
                                (0.002)

tenure                           0.022***
                                (0.003)

Constant                         0.284***
                                (0.104)

-----
Observations                      526
R2                                0.316
Adjusted R2                       0.312
Residual Std. Error              0.441 (df = 522)
F Statistic                       80.391*** (df = 3; 522)
=====
Note:                             *p<0.1; **p<0.05; ***p<0.01
    
```

Motivation

Sampling  
Distributions  
of the OLS  
Estimators

Testing  
Hypotheses  
About a Single  
Population  
Parameter

Testing Against  
One-Sided  
Alternatives

Testing Against  
Two-Sided  
Alternatives

Testing Other  
Hypotheses about  
the  $\beta_j$

Computing  $p$ -Values  
for  $t$  Tests

Practical (Economic)  
versus Statistical  
Significance

Confidence  
Intervals

Testing  
Multiple  
Exclusion

- The correct way to interpret a CI is to remember that the endpoints,  $\hat{\beta}_j - c \cdot se(\hat{\beta}_j)$  and  $\hat{\beta}_j + c \cdot se(\hat{\beta}_j)$ , **change** with each sample (or at least can change).

Endpoints are random outcomes that depend on the data we draw.

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

A 95% CI means is that for 95% of the random samples that we draw from the population,

the interval we compute using the rule  $\hat{\beta}_j \pm c \cdot se(\hat{\beta}_j)$

will include the value  $\beta_j$ .

**But for a particular sample we do not know whether  $\beta_j$  is in the interval.**

- This is similar to the idea that unbiasedness of  $\hat{\beta}_j$  does *not* means that  $\hat{\beta}_j = \beta_j$ . Most of the time  $\hat{\beta}_j$  is not  $\beta_j$ . Unbiasedness means  $E(\hat{\beta}_j) = \beta_j$ .

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- ① Motivation
- ② Sampling Distributions of the OLS Estimators
- ③ Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
  - Computing  $p$ -Values for  $t$  Tests
  - Practical (Economic) versus Statistical Significance
- ④ Confidence Intervals
- ⑤ Testing Multiple Exclusion Restrictions
  - $R$ -Squared Form of the  $F$  Statistic
  - The  $F$  Statistic for Overall Significance of a Regression

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- Sometimes want to test **more than one hypothesis**, which then includes **multiple parameters**.
- Generally, it is not valid to look at individual  $t$  statistics.
- We need a specific statistic used to test **joint hypotheses**.



- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

## Example:

$$\log(\text{wage}) = \beta_0 + \beta_1 \text{educ} + \beta_2 \text{exper} + \beta_3 \text{tenure} + u$$

- Let's consider the following null hypothesis:

$$H_0 : \beta_2 = 0, \beta_3 = 0$$

- **Exclusion Restrictions:** We want to know if we can exclude some variables jointly.

# Testing Multiple Exclusion Restrictions

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- To test  $H_0$ , we need a **joint (multiple) hypotheses test**.
- A  $t$  statistic can be used for a single exclusion restriction; it does not take a stand on the values of the other parameters.
- We are considering the alternative to be:

$$H_1 : H_0 \text{ is not true}$$

- So,  $H_1$  means **at least one** of betas is different from zero.

- The original model, containing all variables, is the **unrestricted model**:

$$\log(\text{wage}) = \beta_0 + \beta_1 \text{educ} + \beta_2 \text{exper} + \beta_3 \text{tenure} + u$$

- When we impose  $H_0 : \beta_2 = 0, \beta_3 = 0$ , we get the **restricted model**:

$$\log(\text{wage}) = \beta_0 + \beta_1 \text{educ} + u$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- We want to see how the fit deteriorates as we remove the two variables.
- We use, initially, the **sum of squared residuals** from the two regressions.
- It is an algebraic fact that the SSR must increase (or, at least not fall) when explanatory variables are dropped. So,

$$SSR_r \geq SSR_{ur}$$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

## *F* test

Does the  $SSR$  increase proportionately by enough to conclude the restrictions under  $H_0$  are false?

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- In the general model:

$$y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + u$$

we want to test that the last  $q$  variables can be excluded:

$$H_0 : \beta_{k-q+1} = 0, \dots, \beta_k = 0$$

- We get  $SSR_{ur}$  from estimating the full model.

# Testing Multiple Exclusion Restrictions

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- The restricted model we estimate to get  $SSR_r$  drops the last  $q$  variables ( $q$  exclusion restrictions):

$$y = \beta_0 + \beta_1 x_1 + \dots + \beta_{k-q} x_{k-q} + u$$

- The **F statistic** uses a degrees of freedom adjustment. In general, we have

$$F = \frac{(SSR_r - SSR_{ur}) / (df_r - df_{ur})}{SSR_{ur} / df_{ur}} = \frac{(SSR_r - SSR_{ur}) / q}{SSR_{ur} / (n - k - 1)}$$

where  $q$  is the number of exclusion restrictions imposed under the null ( $q = 2$  in our example).

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

$$q = \text{numerator df} = df_r - df_{ur}$$

$$n - k - 1 = \text{denominator df} = df_{ur}$$

- The denominator of the  $F$  statistic,  $SSR_{ur}/df_{ur}$ , is the unbiased estimator of  $\sigma^2$  from the unrestricted model.
- Note that  $F \geq 0$ , and  $F > 0$  virtually always holds.
- As a computational device, sometimes the formula

$$F = \frac{(SSR_r - SSR_{ur})}{SSR_{ur}} \cdot \frac{(n - k - 1)}{q}$$

is useful.



# Testing Multiple Exclusion Restrictions

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- Using classical testing, the rejection rule is of the form

$$F > c$$

where  $c$  is an appropriately chosen **critical value**.

## Distribution of $F$ statistic

Under  $H_0$  (the  $q$  exclusion restrictions)

$$F \sim F_{q, n-k-1}$$


i.e., it has an  $F$  distribution with  $(q, n - k - 1)$  degrees of freedom.

- Recall the  $F$  distribution (see slides for **Appendix B**).

# Testing Multiple Exclusion Restrictions

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple

- Suppose  $q = 3$  and  $n - k - 1 = df_{ur} = 60$ . Then the 5% cv is 2.76.



**Question:** Is there a way to compute the  $F$  statistic with the information reported in the standard output from any econometric/statistical package?

- The  $R$ -squared is always reported.
- The SSR is not reported most of the time.
- It turns out that  $F$  tests for exclusion restrictions can be computed entirely from the  $R$ -squareds for the restricted and unrestricted models.
- Notice that,

$$SSR_r = (1 - R_r^2)SST$$

$$SSR_{ur} = (1 - R_{ur}^2)SST$$

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- Therefore,

$$F = \frac{(R_{ur}^2 - R_r^2)/q}{(1 - R_{ur}^2)/(n - k - 1)}$$

- Notice how  $R_{ur}^2$  comes first in the numerator.
- We know  $R_{ur}^2 \geq R_r^2$  so this ensures  $F \geq 0$ .

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

## Example

**unrestricted model:**  $\log(wage) = \beta_0 + \beta_1 educ + \beta_2 exper + \beta_3 tenure + u$

**restricted model:**  $\log(wage) = \beta_0 + \beta_1 educ + u$

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

```

=====
                        Dependent variable:
=====
                        lwage
-----
educ                    0.092***
                        (0.007)

exper                   0.004**
                        (0.002)

tenure                  0.022***
                        (0.003)

Constant                0.284***
                        (0.104)

-----
Observations            526
R2                      0.316
Adjusted R2             0.312
Residual Std. Error    0.441 (df = 522)
F Statistic             80.391*** (df = 3; 522)
=====
Note:                    *p<0.1; **p<0.05; ***p<0.01

```

```

=====
                        Dependent variable:
-----
                        lwage
-----
educ                    0.083***
                        (0.008)

Constant                0.584***
                        (0.097)

-----
Observations            526
R2                      0.186
Adjusted R2             0.184
Residual Std. Error    0.480 (df = 524)
F Statistic             119.582*** (df = 1; 524)
=====
Note:                    *p<0.1; **p<0.05; ***p<0.01

```

Motivation

Sampling Distributions of the OLS Estimators

Testing Hypotheses About a Single Population Parameter

Testing Against One-Sided Alternatives

Testing Against Two-Sided Alternatives

Testing Other Hypotheses about the  $\beta_j$

Computing  $p$ -Values for  $t$  Tests

Practical (Economic) versus Statistical Significance

Confidence Intervals

Testing Multiple Exclusion

- We say that *exper*, and *tenure* are **jointly statistically significant** (or just **jointly significant**), in this case, at any small significance level we want.
- The  $F$  statistic does not allow us to tell which of the population coefficients are different from zero. And the  $t$  statistics do not help much in this example.

# The $F$ Statistic for Overall Significance of a Regression

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

## The $F$ Statistic for Overall Significance of a Regression

- The  $F$  statistic in the **R** output tests a very special null hypothesis.
- In the model:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k + 0$$

the null is that **all slope coefficients are zero**, i.e.,

$$H_0 : \beta_1 = 0, \beta_2 = 0, \dots, \beta_k = 0$$

- This means that none of the  $x_j$  helps explain  $y$ .
- If we cannot reject this null, we have found no factors that explain  $y$ .



# The $F$ Statistic for Overall Significance of a Regression

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- For this test,

$R_r^2 = 0$  (no explanatory variables under  $H_0$ ).

$R_{ur}^2 = R^2$  from the regression.

$$F = \frac{R^2/k}{(1 - R^2)/(n - k - 1)} = \frac{R^2}{(1 - R^2)} \cdot \frac{(n - k - 1)}{k}$$

# The $F$ Statistic for Overall Significance of a Regression

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
- Testing Against One-Sided Alternatives
- Testing Against Two-Sided Alternatives
- Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

- As  $R^2$  increases, so does  $F$ .
- A small  $R^2$  can lead  $F$  to be significant.
- If the  $df = n - k - 1$  is large (because of large  $n$ ),  $F$  can be large even with a “small”  $R^2$ .
- Increasing  $k$  decreases  $F$ .

# The $F$ Statistic for Overall Significance of a Regression

- Motivation
- Sampling Distributions of the OLS Estimators
- Testing Hypotheses About a Single Population Parameter
  - Testing Against One-Sided Alternatives
  - Testing Against Two-Sided Alternatives
  - Testing Other Hypotheses about the  $\beta_j$
- Computing  $p$ -Values for  $t$  Tests
- Practical (Economic) versus Statistical Significance
- Confidence Intervals
- Testing Multiple Exclusion

```

=====
                        Dependent variable:
=====
                        lwage
-----
educ                    0.092***
                        (0.007)

exper                   0.004**
                        (0.002)

tenure                  0.022***
                        (0.003)

Constant                0.284***
                        (0.104)

-----
Observations            526
R2                      0.316
Adjusted R2             0.312
Residual Std. Error    0.441 (df = 522)
F Statistic             80.391*** (df = 3; 522)
=====
Note:                    *p<0.1; **p<0.05; ***p<0.01
    
```

```

=====
                        Dependent variable:
=====
                        lwage
-----
Constant                1.623***
                        (0.023)

-----
Observations            526
R2                      0.000
Adjusted R2             0.000
Residual Std. Error    0.532 (df = 525)
=====
Note:                    *p<0.1; **p<0.05; ***p<0.01
    
```